ABclonal www.abclonal.com

Recombinant Mouse IL-36 gamma/IL-1F9 Protein

Catalog No.: RP01762 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 215257 Q8R460

Tags C-His

Synonyms

If36g; Il1f9; IL-36gamma

Product Information

Source Purification
HEK293 cells > 95% by SDS-

PAGE.

Endotoxin < 0.01EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).

Background

The protein is a member of the interleukin 1 cytokine family. The activity of this cytokine is mediated by interleukin 1 receptor-like 2 (IL1RL2/IL1R-rp2), and is specifically inhibited by interleukin 1 family, member 5 (IL1F5/IL-1 delta). Interferon-gamma, tumor necrosis factor-alpha and interleukin 1, beta (IL1B) are reported to stimulate the expression of this cytokine in keratinocytes. The expression of this cytokine in keratinocytes can also be induced by a contact hypersensitivity reaction or herpes simplex virus infection.

Basic Information

Description

Recombinant Mouse IL-36 gamma/IL-1F9 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly13-Ser164) of mouse IL36G (Accession #) fused with and a 6×His tag at the C-terminus.

Bio-Activity

Measured by its ability to induce IL-6 secretion by NIH \square 3T3 mouse embryonic fibroblast cells. Towne, J.E. et al. (2004) J. Biol. Chem. 279:13677. The ED ₅₀ for this effect is 0.1-0.4 µg/mL,corresponding to a specific activity of $2.5 \times 10 < \sup 3 < \sup 71 \times 10 < \sup 4 < \sup 91$.

Storage


Store at -20° C. Store the lyophilized protein at -20° C to -80° C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20° C for 3 months, at $2-8^{\circ}$ C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com

Validation Data



Recombinant Mouse Interleukin-36 gamma/IL36G Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 20 kDa.



Recombinant Mouse IL36G induce IL-6 secretion by NIH \square 3T3 mouse embryonic fibroblast cells. Towne, J.E. et al. (2004) J. Biol. Chem. 279:13677. The ED $_{50}$ for this effect is 0.1-0.4 μ g/mL,corresponding to a specific activity of $2.5 \times 10^3 \sim 1 \times 10^4$ units/mg.